




EVENING GUIDE · SLEEP

How to read a sleep label slowly on a tired night

For adults who sleep less well than they would like and feel lost among nightstand bottles. You will be able to read any sleep label with care and decide what, if anything, to try.

- **Low and early**
Why small melatonin doses taken hours before bed often make more sense.
- **Label truth**
What serving size, extracts and third party marks actually tell you.
- **One at once**
How to try a single aid with care and know when to stop.

By Tara Whelan · Updated Jul 9, 2026 · 11 min read

CONTENTS

01	Sleep habits before any pill	06	Valerian, chamomile and other herbs
02	How to read any sleep label	07	Interactions, who should pause, and when to get help
03	Quality checks and third party marks	08	The checklist
04	Melatonin for timing, in low doses	09	Sources and further reading
05	Magnesium, glycine, L-theanine and tart cherry		

The lamp is still on and the phone glows beside a small crowd of bottles. Each label promises rest, yet none says plainly what is inside or whether it suits you. Once the words on those labels start to make sense, the night feels kinder and choice grows easier.

01

Sleep habits before any pill

A steady rhythm supports every other choice. Going to bed and rising near the same time helps circadian rhythm stay settled, even on weekends. Light matters through the day. Morning light and dimmer evenings tell the brain when night begins. A short wind down helps too. A warm shower, a book, low voices in the house, and the long exhale before lights out can soften the hour before bed.

Caffeine lingers longer than many expect, so an afternoon cutoff supports easier onset. Alcohol may bring drowsiness at first and then lighter, broken sleep later in the night. The bedroom itself can help. Cool air, hush, dark cloth over small lights, and bedding that feels calm to the touch make rest more likely. Screens in bed do not need shame. Lower brightness, night mode, and putting the phone face down after a set time can be enough.

Persistent trouble deserves attention beyond a bottle. Loud snoring with pauses, gasping, leg movements that disturb sleep, low mood or worry that fills the night, and insomnia that lasts weeks point toward apnea, movement disorders or mental health concerns that need evaluation. A sleep aid cannot fix those patterns on its own. Evidence suggests these habits form the base, and supplements at best support relaxation or timing.

Melatonin is a hormone made by the brain that helps regulate circadian rhythm and signals time to sleep. That signal works best when the rest of the night already leans toward rest.

02

How to read any sleep label

Start with serving size. A gummy, two capsules, or a scoop of powder may each count as one serving, and the amounts listed refer to that serving alone. Then look for amount per serving in milligrams or micrograms. For herbs, look for standardized extract. Only carefully processed roots of *Valeriana officinalis* have been widely studied for sleep, so root and rhizome with a stated marker matter more than a vague leaf powder.

Proprietary blends hide more than they reveal. A blend may list a total weight for five herbs without saying how much of each is present. That makes it hard to judge dose or risk.

Directions and warnings deserve the same slow reading. Slow release versus standard changes timing. In the UK, short term insomnia in adults is usually treated with one 2 mg slow release tablet taken 1 to 2 hours before bedtime for 1 to 4 weeks and up to 13 weeks. The pattern is short courses rather than open ended nightly use.

Natural does not mean safe. Valerian can cause headache, dizziness, stomach problems or sleeplessness. Tart cherry juice contains quercetin, which can interact with certain blood thinners and anticoagulants. In the US, sleep supplements are classified as dietary supplements under the Dietary Supplement Health and Education Act of 1994 and can be sold without FDA preapproval of safety or efficacy. The label is written by the seller, not a neutral referee.

FROM THE EDITORS

Third party testing checks identity, strength and contaminants, but it does not prove that a product helps sleep. Look for USP, NSF or ConsumerLab marks, plus lot numbers and expiration dates. Avoid blends that hide doses and labels that promise sleep outright.

03

Quality checks and third party marks

A careful label invites checking. Third party marks from USP, NSF or ConsumerLab suggest an outside lab has looked at what is in the bottle. They check whether the contents match the label and whether unwanted substances are present. Lot numbers and expiration dates allow a batch to be traced. Clear amounts for each ingredient allow a fair comparison from one product to another.

Red flags are common at dusk in the supplement aisle. High doses of melatonin, missing amounts in a blend, claims of a perfect night, and no mention of side effects or interactions suggest caution. Product variability is well documented. A study of 31 melatonin supplements

found real content varied from minus 83 percent to plus 478 percent of labeled concentration. In a 2023 JAMA study of 25 melatonin gummy products, 22 products were inaccurately labeled with actual quantity from 74 to 347 percent of label.

Absorption adds more uncertainty. Melatonin bioavailability has a broad range of 1 percent to 74 percent due to dose and supplement formulation. What is swallowed is not always what reaches the blood. For this reason, a low labeled dose may act differently from one person to another and from one brand to another. Choosing a tested product with a clear single ingredient list lowers, but does not remove, that uncertainty.

04

Melatonin for timing, in low doses

Melatonin suits timing questions more than general sleeplessness. It signals night to the body rather than forcing sedation. Low melatonin doses of 0.3 mg and 1 mg were observed to stimulate night levels like those in young adults. As the UC Davis Health article on melatonin puts it, "Small doses can be effective. It's not the case that the more melatonin you take, the more effective it will be, or the sleepier you will be."

Timing matters as much as amount. For best rhythm effects, take melatonin roughly 3 to 4 hours before desired sleep time, such as 6 or 7 PM for an 11 PM bedtime. That early hour can feel strange at first. It reflects how long the signal needs to shift rhythm, which differs from taking a pill at lights out. For jet lag in adults, the usual UK dose is one 3 mg standard tablet at normal bedtime at destination, up to 6 mg, for a maximum of 5 days. Short use fits travel rather than months of nightly tablets.

Side effects tend to be mild but real. Common side effects of melatonin include headache and drowsiness, reported in 1 in 10 people in package information. Daytime grogginess, vivid dreams and stomach upset are also reported. Long term safety of nightly melatonin use beyond one to two months is not established, so evidence does not show that continuous use stays safe or stays helpful.

Some people need medical supervision before any trial. Melatonin should not be combined with fluvoxamine or tyrosine kinase inhibitors, and people with epilepsy or on blood thinners need medical supervision. Those cautions underline why one sleep aid at a time, in a low dose, for a short period, tells a clearer story than a mix of bottles.

05

Magnesium, glycine, L-theanine and tart cherry

Magnesium appears in many nightstand formulas, yet evidence for sleep is thin and mostly linked to deficiency, leg cramps or restless legs. People at greatest risk of magnesium deficiency include women, older adults, regular alcohol drinkers, and those with diets high in processed foods. Proton pump inhibitors and diuretics can contribute to magnesium deficiency by impairing absorption or increasing urinary loss. Whether correcting levels helps sleep in people without deficiency remains unclear.

Forms differ in feel and cost. Most evidence for magnesium as a sleep aid is for magnesium citrate, but it has potent laxative effects, while glycinate is gentler and oxide is less expensive. Mayo Clinic source recommends 250 to 500 milligrams of magnesium in a single dose at bedtime for people with healthy kidney function. Those with kidney disease face higher risk and need clinician advice before use. Oral magnesium is recommended over topical sprays or gels because transdermal absorption is low or inefficient. Denise M. Millstine, M.D., director of integrative medicine and health at Mayo Clinic in Arizona, says, "I wouldn't recommend topical magnesium, whether it's a spray or a gel,".

Glycine and L-theanine work through quieter ideas. Glycine may regulate sleep wake cycles when taken before bed by helping lower body temperature, with efficacy even at small doses. The cooling idea fits the natural drop in core temperature before sleep. L-theanine from tea leaves is postulated to promote relaxation and may reduce night wakings without causing daytime sleepiness. Trials are small or short, so exact best dose and duration for glycine and L-theanine for sleep are not established.

Tart cherry brings food as much as supplement. Tart cherries contain melatonin, anthocyanins, serotonin and tryptophan with antioxidant and anti-inflammatory effects that may support sleep. A systematic review of seven interventional studies found three reported better sleep duration, efficiency or onset after tart cherry, and three reported higher melatonin levels. Studies are few, small and varied in dose, duration and population, so clinical benefit is not proven. Juice adds sugar and fluid before bed, which some bladders may notice.

06

Valerian, chamomile and other herbs

Valerian carries a long tradition and a mixed record. Valerian seems most effective after regular use for two or more weeks, but dosage varied across studies and optimal dose is unclear. The idea often involves GABA, a calming brain signal, though the mechanism remains uncertain. A 2024 literature review in *Psychiatry Investigation* concluded valerian, hops and melatonin could improve sleep quality but strength of evidence varies and optimal dosages and durations are unestablished.

Valerian trials are mixed and inconclusive, some people feel more awake, and optimal dose and duration are not established. Daytime drowsiness can follow nighttime use. Valerian should not be taken with other sedatives including benzodiazepines, central nervous system depressants and barbiturates. It should be avoided with liver disease and in pregnancy or breastfeeding. Slow patience suits this herb better than a single night test, yet patience should still include a stop date if nothing changes.

Chamomile, hops and lavender live mostly in tradition and small trials. Warm tea before bed may soothe as a ritual even when proof for sleep remains limited. Traditional use tells us what people hoped for across generations. Modern evidence asks a narrower question about dose, duration and measured sleep, and often finds too little to judge. That gap does not make a cup of chamomile wrong. It means claims of strength should stay modest.

Comparison can steady choice when shelves feel crowded.

AID	WHAT IT MAY OFFER	WHAT TO KEEP IN MIND
Melatonin	Timing support for rhythm and jet lag	Low doses, early timing, short courses, label variability
Magnesium	Support where intake is low	Citrate may loosen bowels, kidney disease needs care, sprays absorb poorly
Glycine	Possible cooling before sleep even at small doses	Best dose and duration not established
L-theanine	Relaxation with less chance of daytime sleepiness	Trials small, effects on wakings vary
Tart cherry	Food source with melatonin and anthocyanins	Few small studies, sugar and fluid, quercetin and blood thinner caution
Valerian	Possible sleep quality support after two or more weeks	Mixed trials, daytime drowsiness, avoid with sedatives and liver disease

FROM THE EDITORS

Try one aid at a time so effects and side effects stay clear. Keep a simple note of dose, timing and morning feel for one to two weeks. Stop and seek care for chest pain, trouble breathing, severe daytime sleepiness, signs of allergy, or sleep that keeps worsening.

Interactions, who should pause, and when to get help

Combinations create most of the risk at night. Blood thinners, SSRIs, sedatives and alcohol, blood pressure and diabetes drugs, antibiotics, diuretics, transplant and seizure history all call for a pharmacist or clinician check before any trial. Valerian adds sedation when mixed with other sedatives. Quercetin in tart cherry can affect certain blood thinners and anticoagulants. Magnesium can interact with antibiotics and diuretics, and those medicines can also shift magnesium balance.

Some groups should pause altogether unless a clinician advises otherwise. Pregnancy, breastfeeding, liver and kidney disease, and childhood need extra caution. Children using melatonin need pediatric guidance rather than adult dosing. Older adults may feel stronger effects from drowsiness and should start with added care around falls and morning driving.

Knowing when to stop matters as much as knowing how to start. Stop a trial if headaches, dizziness, nausea, daytime sleepiness, low mood, or vivid agitation continue. Seek care sooner for signs of allergy, irregular heartbeat, fainting, or thoughts that frighten you. Bring bottles to the visit, with doses and timing written down. A short note from the nightstand often helps the clinician more than a tracker score.

The checklist

-
- ☐ Keep a steady bedtime and rise time for two weeks first
 - ☐ Dim lights and lower screens in the hour before bed
 - ☐ Set an afternoon caffeine cutoff and note alcohol effects
 - ☐ Read serving size and amount per serving before price
 - ☐ Choose single ingredients over blends that hide doses
 - ☐ Look for USP, NSF or ConsumerLab marks and lot numbers
 - ☐ Try one aid at a time at the lowest studied amount
 - ☐ Take melatonin hours before bed when using it for timing
 - ☐ Check interactions with blood thinners, sedatives and antibiotics
 - ☐ Pause and ask a clinician if pregnant, breastfeeding or chronically ill
 - ☐ Stop if side effects linger or sleep keeps worsening
 - ☐ Bring bottles and notes to any care visit
-

Sources and further reading

- Melatonin dose and timing: Do we have it right? · [pmc.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)
- Magnesium for sleep: What you need to know about its benefits · mcpress.mayoclinic.org
- The Effect of Tart Cherry on Sleep Quality and Sleep Disorders: A Systematic Review · [pmc.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)
- The Science Behind Sleep Supplements: Clinical Guidance Amidst Regulatory Concerns · pulmonologyadvisor.com
- How and when to take melatonin · nhs.uk
- Herbal and Natural Supplements for Improving Sleep: A Literature Review · psychiatryinvestigation.org
- 7 Sleep Aids That Don't Mix With Prescription Drugs · aarp.org
- Valerian for insomnia? Maybe-Valerian: A safe and effective herbal sleep aid? · mayoclinic.org

For information only, not medical advice. Ongoing trouble sleeping, loud snoring with pauses in breathing, or low mood that lingers deserves a doctor's attention, and supplements can interact with medication.



WRITTEN BY

Tara Whelan

Tara runs the night desk and the evening guides at Well with Luna. She has tested every mask, nasal strip and pillowcase we have written about, one night at a time, and keeps her notes in a paper diary by the bed.
